

Atom lens without chromatic aberrations

Maxim A. Efremov,^{1,2} Polina V. Mironova,¹ Wolfgang P. Schleich¹

¹*Institut für Quantenphysik and Center for Integrated Quantum Science and Technology (IQST), Universität Ulm, 89081 Ulm, Germany*

²*A.M. Prokhorov General Physics Institute, Russian Academy of Sciences, 119991 Moscow, Russia**

(Dated: December 3, 2012)

We propose a lens for atoms with reduced chromatic aberrations and calculate its focal length and spot size. In our scheme a two-level atom interacts with a near-resonant standing light wave formed by two running waves of slightly different wave vectors, and a far-detuned running wave propagating perpendicular to the standing wave. We show that within the Raman-Nath approximation and for an adiabatically slow atom-light interaction, the phase acquired by the atom is independent of the incident atomic velocity.

A crucial element of the tool box of atom optics [1, 2] is a lens to focus atom waves. Many suggestions have been given [3] and realized in experiments [4]. However, most of these realizations suffer from chromatic aberrations. In the present paper we propose a lens, which is free of this type of aberrations using a combination of light waves, which introduce two different time scales into the interaction of the atom with the fields. Our proposal can be implemented with today's technology.

In order to compare and contrast the features of our proposed lens with the corresponding ones of a conventional lens, we briefly rederive the familiar result [5] that the focal length and spot size of a thin lens created by an optical potential depend quadratically and linearly on the velocity of the atom, respectively. In contrast, for our lens the focal length depends linearly on and the spot size is independent of the velocity. In our scheme we scatter a two-level atom from a standing light wave built out of two running waves propagating in almost antiparallel directions. The atom moves with a large velocity orthogonal to the standing wave and experiences a Doppler shift as it traverses the field. Moreover, in order to control the sign of the effective detuning, we use a running wave with an envelope in “top-hat” shape, which is strongly detuned from the atomic transition and propagates orthogonal to the standing wave.

Within the adiabatic approximation and the Raman-Nath approximation [6] for the center-of-mass motion we find the wave function of the ground state accumulates a position-dependent phase. Near the nodes of the standing wave this phase depends quadratically on the position of the atom as in a conventional lens. However, now the focal length and the spot size are multiplied by the ratio of the detuning and the Doppler shift. Since the latter depends on the velocity of the atom, in our lens the velocity dependence of the focal length and the spot size is reduced by one power, resulting in a lens without chromatic aberrations.

We start our analysis by recalling the key features of a conventional lens where a two-level atom interacts with a standing light field detuned by Δ giving rise to the Rabi frequency Ω_0 . This interaction creates the optical poten-

tial $U_{\text{opt}}(x) \equiv (\hbar\Omega_0^2/\Delta) \sin^2(k_x x)$ for the motion along the x -axis, which we treat quantum-mechanically. For the interaction time $\tau \equiv w_0/v_y$ determined by the waist w_0 of the standing wave and the longitudinal velocity v_y of the atom we imprint the phase

$$\phi(x) \sim -\frac{U_{\text{opt}}(x)\tau}{\hbar} = -\frac{\Omega_0^2 w_0}{\Delta v_y} \sin^2(k_x x) \quad (1)$$

onto the wave function of the center-of mass motion of the atom in the ground state.

Within the parabolic approximation, $k_x|x| \ll 1$, the phase

$$\phi(x) \approx -\frac{\Omega_0^2 w_0}{\Delta v_y} k_x^2 x^2 \quad (2)$$

is a quadratic function of x . Since the transverse momentum p of the atom of the mass M follows from the derivative of this phase, we find that

$$p \equiv Mv \equiv \hbar \frac{\partial \phi}{\partial x} = -2 \frac{\hbar\Omega_0^2 w_0}{\Delta} k_x^2 x \quad (3)$$

depends linearly on x .

The focal length $\mathcal{F}_0$ of a lens is defined [5, 6] as the distance traveled by the atom in the longitudinal direction during the time necessary to go with the momentum p from the position x to $x = 0$, which yields

$$\mathcal{F}_0 = -v_y \frac{x}{(p/M)} = \frac{Mv_y^2}{2\hbar\Omega_0} \frac{\Delta}{k_x^2 w_0 \Omega_0}. \quad (4)$$

The spot size $\mathcal{S} \equiv \alpha_0 \mathcal{F}$ is determined by the focal length $\mathcal{F}$ and the angular divergence $\alpha_0 \equiv \delta v/v_y$ of the atomic beam, with δv being the uncertainty of the transverse atomic velocity. For a Gaussian wave packet of width δx the uncertainty $\delta v = \hbar/(M\delta x)$ gives rise to the angular divergence $\alpha_0 = \hbar/(Mv_y\delta x)$ and the spot size

$$\mathcal{S}_0 = \frac{v_y \Delta}{2k_x^2 \delta x w_0 \Omega_0^2}. \quad (5)$$

According to Eqs. (4) and (5) in a thin conventional lens both the focal length $\mathcal{F}_0$ and the spot size $\mathcal{S}_0$ depend on the atomic velocity v_y , namely $\mathcal{F}_0 \propto v_y^2$ and




Figure 1: Scattering of the wave packet $\Psi = \Psi(x)$ of a two-level atom by a combination of a standing electromagnetic field formed by two propagating waves of wave vectors $\mathbf{k}_1$ and $\mathbf{k}_2$ and a traveling wave propagating orthogonal to the $x-y$ -plane serving as a control field. As the atom propagates along the y -axis with the velocity $\mathbf{v} = v_y \mathbf{e}_y$, the envelope of the two running waves translates according to the relation $y = v_y t$ into the time-dependent function $f_l = f_l(v_y t)$. During the atom-field interaction the effective detuning of the field frequencies from the atomic transition changes its sign due to the control field with an envelope $f_c(t)$ in the shape of a “top-hat”.

$S_0 \propto v_y$, resulting in large chromatic aberrations. These scaling laws serve as our motivation to engineer a phase element for atoms and, in particular, a lens with reduced chromatic aberrations. Our suggestion relies on the interaction of two-level atom with a near-resonant standing wave providing us with the optical potential inducing the focusing, and a far-detuned traveling light wave removing the achromatic aberrations.

We create the lens field by the superposition

$$\mathbf{E}_s(t, \mathbf{r}) = \mathbf{E}_l(\mathbf{r})e^{-i\omega t} (e^{i\mathbf{k}_2 \mathbf{r}} - e^{i\mathbf{k}_1 \mathbf{r}}) + c.c. \quad (6)$$

of two traveling waves of wave vectors $\mathbf{k}_1 \equiv (k_x, k_y) \equiv (k \cos \alpha, k \sin \alpha)$ and $\mathbf{k}_2 \equiv (-k_x, k_y)$, which form an angle α relative to the x -axis shown in Fig. 1. Here $\mathbf{E}_l(\mathbf{r})$ describes the position-dependent real-valued amplitude of the waves, which, throughout the article, is assumed to be of the form of the TEM₀₁ Hermite-Gauss mode with a node along the y -axis. The frequency ω is detuned from the frequency of the atomic transition between the ground $|g\rangle$ and excited $|e\rangle$ states of the corresponding energies $E_g \equiv \hbar\omega_g$ and $E_e \equiv \hbar\omega_e$ by an amount $\Delta \equiv \omega_e - \omega_g - \omega$ as shown in Fig. 1.

A running control wave

$$\mathbf{E}_r(t, \mathbf{r}) = \mathbf{E}_c(x, y)e^{i(k_c z - \omega_c t)} + c.c. \quad (7)$$

with the position-dependent amplitude $\mathbf{E}_c(x, y)$ and the

shape of a “top-hat” propagates along the z -axis perpendicular to the xy -plane. The frequency ω_c is far-detuned by $\Delta_c \equiv \omega_{e'} - \omega_e - \omega_c$ from the atomic transition between the excited state $|e\rangle$ and some other state $|e'\rangle$. We suppose that the control field is weak enough to be considered perturbatively, resulting in the Stark shift $\Delta E_e = -|\tilde{\boldsymbol{\varphi}} \mathbf{E}_c|^2 / (\hbar \Delta_c)$ of the atomic excited state $|e\rangle$, where $\tilde{\boldsymbol{\varphi}} \equiv \langle e | \mathbf{d} | e' \rangle$ is the dipole matrix element.

Throughout the article we assume that the velocity v_y of the atom in the direction of the y -axis is large and remains almost constant during the scattering process. For this reason we consider this motion classically, which allows us to set $y \equiv v_y t$. In contrast, the motion along the x -axis is described quantum mechanically within the Raman-Nath approximation [5, 6], that is the displacement of the atom along the x -axis caused by its interaction with the standing light wave is assumed to be small compared to its period. Moreover, due to the small interaction time and the large detuning Δ we neglect the spontaneous emission of atom in the excited state.

In the framework of these approximations the time evolution of the state-vector

$$|\Psi(t)\rangle = a_e(t; \mathbf{r})e^{-i\omega_e t}|e\rangle + a_g(t; \mathbf{r})e^{-i\omega_g t}|g\rangle \quad (8)$$

follows from the Schrödinger equation. Indeed, within the rotating-wave approximation the time-dependent amplitudes a_g and a_e , which depend on the position $\mathbf{r}$ of the atom as a parameter, obey the system of equations

$$i\hbar \frac{d}{dt} \begin{pmatrix} a_e \\ a_g \end{pmatrix} = \hat{H} \begin{pmatrix} a_e \\ a_g \end{pmatrix}. \quad (9)$$

The Hamiltonian

$$\hat{H} \equiv \begin{pmatrix} \Delta E_e & V_l^* e^{i\Delta t} \\ V_l e^{-i\Delta t} & 0 \end{pmatrix} \quad (10)$$

contains the complex-valued coupling matrix elements

$$V_l(\mathbf{r}) = 2\boldsymbol{\varphi} \mathbf{E}_l(x, y)e^{-ik_y y} \sin(k_x x), \quad (11)$$

with $\boldsymbol{\varphi} \equiv \langle g | \mathbf{d} | e \rangle$ being the dipole matrix element.

We assume that the x - and y -dependence of $\mathbf{E}_l$ and $\mathbf{E}_c$ can be separated and, since the atomic motion along the y -axis is treated classically, $y = v_y t$, we find the form $\mathbf{E}_l(x, y) \equiv \mathcal{E}_l(x)f_l(y) = \mathcal{E}_l(x)f_l(v_y t)$ and $\mathbf{E}_c(x, v_y t) = \mathcal{E}_c(x)f_c(t)$ for the electric field amplitudes. Here the envelope function

$$f_l(y) \equiv \frac{\sqrt{2}}{\pi^{1/4} w_0} \exp\left(-\frac{y^2}{2w_0^2}\right) \quad (12)$$

of the standing wave results from the TEM₀₁ Hermite-Gauss mode along the y -axis and satisfies the normalization conditions

$$\int_{-\infty}^{\infty} dy f_l^2(y) = w_0. \quad (13)$$

In contrast, the envelope

$$f_c(t) \equiv \theta(t) = \begin{cases} 1, & t \geq 0 \\ 0, & t < 0 \end{cases} \quad (14)$$

of the control field has a "top-hat" profile, that is a step-wise dependence as expressed by the Heaviside function $\theta(t)$ [7].

Moreover, the detunings Δ and Δ_c are assumed to have the same sign and we can then choose the amplitude of the Rabi frequency $\Omega_c \equiv |\tilde{\rho}\mathcal{E}_c|/\hbar$ of the control field as

$$\Omega_c = \sqrt{2\Delta\Delta_c}. \quad (15)$$

As a result, the Stark shift ΔE_e induced by the control light field is given by $\Delta E_e = -2\hbar\Delta f_c^2(t)$.

The Hamiltonian Eq. (10) takes finally the form

$$\hat{H} \cong \begin{pmatrix} -2\hbar\Delta f_c^2(t) & \hbar\Omega(x)f_l(v_y t)e^{i(\Delta+\omega_\alpha)t} \\ \hbar\Omega(x)f_l(v_y t)e^{-i(\Delta+\omega_\alpha)t} & 0 \end{pmatrix}, \quad (16)$$

where

$$\omega_\alpha \equiv k_y v_y \quad (17)$$

and

$$\Omega(x) \equiv (2|\tilde{\rho}\mathcal{E}_l|/\hbar) \sin(k_x x) \equiv \Omega_0 \sin(k_x x) \quad (18)$$

are the velocity-dependent Doppler and position-dependent Rabi frequencies, respectively.

We now solve the Schrödinger equation (9) with the Hamiltonian Eq. (16) in the case of the adiabatically slow atom-field interaction. For this purpose we substitute the second equation of system Eq. (9) for the amplitude a_g into the first one for a_e and get the second order differential equation

$$\frac{d^2}{dt^2}a_g + \left(i\tilde{\Delta} - \frac{1}{f_l} \frac{df_l}{dt}\right) \frac{d}{dt}a_g + \Omega^2 f_l^2 a_g = 0 \quad (19)$$

with the initial conditions

$$a_g(t_0) = 1, \quad \left. \frac{da_g}{dt} \right|_{t_0} = 0 \quad (20)$$

at time t_0 . Here we have introduced the time-dependent effective detuning

$$\tilde{\Delta}(t) \equiv \Delta + \omega_\alpha - 2\Delta f_c^2(t) = \Delta + \omega_\alpha - 2\Delta\theta(t). \quad (21)$$

In the case of a slowly varying envelope $f_l(t)$ with

$$\left| \frac{1}{f_l} \frac{df_l}{dt} \right| \ll |\tilde{\Delta}|, \quad \text{or} \quad |\tilde{\Delta}|\tau \gg 1, \quad (22)$$

we can neglect its time derivative in the second term of Eq. (19) and arrive at the approximate equation

$$\frac{d^2}{dt^2}a_g + i\tilde{\Delta} \frac{d}{dt}a_g + \Omega^2 f_l^2 a_g \simeq 0. \quad (23)$$

For each time interval, that is for $-\infty < t \leq 0$ and $0 \leq t < \infty$, when the detuning $\tilde{\Delta}$ is constant, the solution of Eq. (23) with the initial equation Eq. (20) reads

$$a_g(t) = \frac{1}{2} \left[1 + \frac{\tilde{\Delta}(t_0)}{\lambda(t_0)} \right] e^{i\phi_+(t)} + \frac{1}{2} \left[1 - \frac{\tilde{\Delta}(t_0)}{\lambda(t_0)} \right] e^{i\phi_-(t)} \quad (24)$$

with

$$\phi_\pm(t) = \frac{1}{2} \int_{t_0}^t dt' [\pm\lambda(t') - \tilde{\Delta}(t')] \quad (25)$$

and

$$\lambda(t) \equiv \sqrt{\tilde{\Delta}^2(t) + 4\Omega^2 f_l^2(v_y t)}. \quad (26)$$

For the two time intervals $-\infty < t \leq 0$ and $0 \leq t < \infty$ the initial time t_0 corresponds to $t_0 = -\infty$ and $t_0 = 0$, respectively, where the envelope $f_l(v_y t)$ vanishes. From Eq. (26) we find $\lambda(t_0) = |\tilde{\Delta}(t_0)|$ and

$$\frac{1}{2} \left[1 \pm \frac{\tilde{\Delta}(t_0)}{\lambda(t_0)} \right] = \frac{1}{2} \left[1 \pm \frac{\tilde{\Delta}(t_0)}{|\tilde{\Delta}(t_0)|} \right] = \theta[\pm\tilde{\Delta}(t_0)]. \quad (27)$$

With the definitions Eqs. (21), (25) and (26) of $\tilde{\Delta}$, $\phi_\pm$ and λ together with the explicit form Eq. (27) of the initial condition we can cast Eq. (24) into the compact form $a_g \equiv \exp(i\varphi_g)$, where the phase

$$\varphi_g(t; x) = \frac{1}{2} \int_{-\infty}^t dt' \left[\text{sgn}(\tilde{\Delta}) \sqrt{\tilde{\Delta}^2 + 4\Omega^2(x) f_l^2(v_y t')} - \tilde{\Delta} \right] \quad (28)$$

depends on the transverse coordinate x of the atom. Since we are interesting in engineering a lens for matter waves, we can ignore the phase $-\omega_g t$ in the Schrödinger picture, which is independent of x , and the total phase Φ_g acquired by the atom during its interaction with the two light fields is the sum

$$\Phi_g \equiv \varphi_g(t \rightarrow \infty; x) = \phi_g(x; \Delta + \omega_\alpha) + \phi_g(x; -\Delta + \omega_\alpha) \quad (29)$$

of the two contributions determined by the time intervals $-\infty < t \leq 0$ and $0 \leq t < \infty$, where

$$\phi_g(x; \delta) \equiv \frac{\text{sgn}(\delta)}{2} \int_0^\infty dt \left[\sqrt{\delta^2 + 4\Omega^2(x) f_l^2(v_y t)} - |\delta| \right]. \quad (30)$$

In the case of $\omega_\alpha \ll |\Delta|$, the total phase Φ_g given by Eq. (29) reduces to

$$\Phi_g(x) = \omega_\alpha \int_0^\infty dt \left[\frac{|\Delta|}{\sqrt{\Delta^2 + 4\Omega^2(x) f_l^2(v_y t)}} - 1 \right]. \quad (31)$$

When we introduce the integration variable $y \equiv v_y t$, and recall the definitions Eqs. (17) and (18) of ω_α and

Ω we arrive at

$$\Phi_g(x) = k_y \int_0^\infty dy \left\{ \left[1 + 4 \frac{\Omega_0^2}{\Delta^2} \sin^2(k_x x) f_l^2(y) \right]^{-\frac{1}{2}} - 1 \right\}. \quad (32)$$

We emphasize that Φ_g is proportional to k_y , which is a consequence of the non-collinearity of the two wave vectors $\mathbf{k}_1$ and $\mathbf{k}_2$. Moreover, Φ_g is independent of v_y . Hence, the combination of the lens field and the control wave acting on the atom creates an achromatic phase element.

We now use this phase element to construct a lens with reduced chromatic aberrations. For this purpose we consider a position of the atom close to a node of the standing wave, which allows us to expand the square root in Eq. (32), and we arrive at

$$\Phi_g(x) \approx \frac{k_y v_y}{\Delta} \left(-\frac{\Omega_0^2}{\Delta} \frac{w_0}{v_y} k_x^2 x^2 \right) = \frac{k_y v_y}{\Delta} \phi(x). \quad (33)$$

Here we have recalled the normalization condition Eq. (13) for the profile function f_l and the form Eq. (2) of the phase ϕ induced by a regular optical potential.

Due to the control field Φ_g is the product of the phase ϕ corresponding to a conventional optical potential and the ratio $(k_y v_y)/\Delta$. Hence, the focal length and the spot size of our lens read

$$\mathcal{F} = \frac{\Delta}{k_y v_y} \mathcal{F}_0 \quad (34)$$

and

$$\mathcal{S} = \frac{\Delta}{k_y v_y} \mathcal{S}_0. \quad (35)$$

Since according to Eqs. (4) and (5) $\mathcal{F}_0$ and $\mathcal{S}_0$ depend quadratically and linearly on v_y , in our lens the focal length $\mathcal{F}$ is proportional to v_y and the spot size $\mathcal{S}$ is independent of it. This scaling implies a reduction of the chromatic aberrations in comparison with the conventional technique of focusing atoms. Moreover, in our lens both the focal length $\mathcal{F}$ and the spot size $\mathcal{S}$ are larger by a factor $|\Delta|/(k_y v_y)$ than those of the conventional lens.

In summary we have proposed a lens with reduced chromatic aberrations. Our scheme differs from a conventional lens by the use of *two* rather than a single light field. The improvement factor is given by the ratio of the detuning and a Doppler shift. Our calculations have relied on the adiabatic approach using the dressed state picture.

It is interesting to note that we could have also obtained the phase Φ_g given by Eq. (32) by considerations familiar from the Berry phase [8–10]. In this way Φ_g could be interpreted as a sum of two Berry phases acquired by the atom while entering and leaving the fields [11]. Since the Berry phase is purely of geometrical nature, it is insensitive to small perturbations in the control parameters [12]. For this reason, our proposed lens

is more robust against small fluctuations in the system parameters, such as the intensity of the light fields.

Acknowledgments. We are deeply indebted to J. Baudon, M.V. Fedorov, R. Kaiser, M.K. Oberthaler, R. Walser, and V.P. Yakovlev for many suggestions and stimulating discussions. MAE is grateful to the Alexander von Humboldt Stiftung and Russian Foundation for Basic Research (grant 10-02-00914-a). PVM acknowledges support from the EU project "CONQUEST" and the German Academic Exchange Service (DAAD).

* Electronic address: max.efremov@gmail.com

- [1] A.D. Cronin, J. Schmiedmayer, and D.E. Pritchard, *Rev. Mod. Phys.* **81**, 1051 (2009); C.S. Adams, M. Sigel, and J. Mlynek, *Phys. Rep.* **240**, 143 (1994)
- [2] P.R. Berman, Ed., *Atom Interferometry* (Academic Press, San Diego, 1997)
- [3] V.I. Balykin and V.S. Letokhov, *Opt. Commun.* **64**, 151 (1987); G.M. Gallatin and P.L. Gould, *J. Opt. Soc. Am. B* **8**, 502 (1991); B. Dubetsky and P.R. Berman, *Phys. Rev. A* **58**, 2413 (1998); J.L. Cohen, B. Dubetsky, and P.R. Berman, *Phys. Rev. A* **60**, 4886 (1999); S. Meneghini, V.I. Savichev, K.A.H. van Leeuwen and W.P. Schleich, *Applied Physics B* **70**, 675 (2000); V.I. Balykin and V.G. Minogin, *Phys. Rev. A* **77**, 013601 (2008);
- [4] O. Carnal, M. Sigel, T. Sleator, H. Takuma, and J. Mlynek, *Phys. Rev. Lett.* **67**, 3231 (1991); B. Holst and W. Allison, *Nature* **390**, 244 (1997); W.G. Kaenders, F. Lison, A. Richter, R. Wynands, and D. Meschede, *Nature* **375**, 214 (1995); J.J. Berkhout, O.J. Luiten, I.D. Setija, T.W. Hijmans, and T. Mizusaki, *Phys. Rev. Lett.* **63**, 1689 (1989); J.E. Bjorkholm, R.R. Freeman, A. Ashkin, and D.B. Pearson, *Phys. Rev. Lett.* **41**, 1361 (1978); T. Sleator, T. Pfau, V. Balykin and J. Mlynek, *Applied Physics B* **54**, 375 (1992); G. Timp, R.E. Behringer, D.M. Tennant, J.E. Cunningham, M. Prentiss, and K.K. Berggren, *Phys. Rev. Lett.* **69**, 1636 (1992); J.J. McClelland, R.E. Scholten, E.C. Palm, and J. Celotta, *Science* **262**, 877 (1993); M. Mützel, D. Haubrich, and D. Meschede, *Applied Physics B* **70**, 689 (2000); W.H. Oskay, D.A. Steck, and M.G. Raizen, *Phys. Rev. Lett.* **89**, 283001 (2002)
- [5] W.P. Schleich, *Quantum Optics in Phase Space*, (Wiley-VCH, Weinheim, 2001)
- [6] A.P. Kazantsev, G.I. Surdutovich, and V.P. Yakovlev, *Mechanical Action of Light on Atoms* (World Scientific, Singapore, 1990)
- [7] In practice the "top-hat" profile is modeled by a continuous function, for instance, by $\{1 + \tanh(t/\tau_1)\}/2$ with the duration time τ_1 . However, if $\tau_1 \ll \tau$, results obtained with the function $f_c(t)$ Eq. (14) are similar to those obtained with the continuous function.
- [8] M.V. Berry, *Proc. R. Soc. London A* **392**, 45 (1984)
- [9] A. Shapere and F. Wilczek, *Geometric Phases in Physics* (World Scientific, Singapore, 1989)
- [10] A. Bohm, A. Mostafazadeh, H. Koizumi, Q. Niu, and J. Zwanziger, *The geometric phase in quantum systems. Foundations, mathematical concepts, and applications in molecular and condensed matter physics* (Springer,

- Berlin, Heidelberg, 2003)
- [11] P.V. Mironova, M.A. Efremov, and W.P. Schleich, Phys. Rev. A (to be submitted); P.V. Mironova, PhD Thesis at the University of Ulm (2011)
- [12] G. De Chiara and G.M. Palma, Phys. Rev. Lett. **91**, 090404 (2003)